

All Databases
Search PubMed

PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books
for: [Go] [Clear]

Limits Preview/Index History Clipboard Details
Display Abstract [Show] 20 [Sort by] [Send to]
All: 1 Review: 0

About Entrez

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
My NCBI (Cubby)

Related Resources

Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

1: IUBMB Life. 2002 Mar;53(3):161-6.

Related Articles, Links

LAST motifs and SMART domains in gene 32 protein: an unfolding story of autoregulation?

Karpel RL.

Department of Chemistry and Biochemistry, University of Maryland, Baltimore County, Baltimore 21250, USA. karpel@umbc.edu

Bacteriophage T4 gene 32 protein is a classical single strand-specific DNA binding protein. It is a single polypeptide chain of 301 amino acid residues that consists of three structural domains, each of which has a binding function. The N-terminal domain is involved in homotypic protein-protein interaction (the basis of binding cooperativity), the core domain binds single strands directly, and the C-terminal domain has a role in heterotypic protein-protein association. The three domains have traditionally been thought to be independent of each other. However, the observation of a striking repetition of a basic, polar sequence (the "LAST" Motif), seen in both the N-terminal and core domains, suggests a linkage between these domains. Moreover, the C-domain and adjoining portion (flap) of the core are highly acidic, and are potential mimics of single-stranded DNA. With these observations, I construct a model in which this flap is associated with the ssDNA binding site in the absence of DNA, and upon cooperative protein binding to DNA, the flap now associates with the N-terminal domain of the adjacent DNA-bound protein. The flap thus acts as a gate, which might slow the binding of the protein to DNA. This could lead to the regulation of the protein's various interactions with other proteins, as well as affect its ability to lower DNA melting temperature.

PMID: 12102172 [PubMed - indexed for MEDLINE]

Display Abstract [Show] 20 [Sort by] [Send to]


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

May 29 2005 17:30:14



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for

Go Clear

Limits

Preview/Index

History

Clipboard

Details

Display

Abstract

Show 20

Sort by

Send to

All: 1

Review: 0

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Mol Microbiol. 2005 Mar;55(5):1502-14.

Related Articles, Links

Protein-DNA interactions in the T4 dNTP synthetase complex dependent on gene 32 single-stranded DNA-binding protein.

Kim J, Wheeler LJ, Shen R, Mathews CK.

Department of Biochemistry and Biophysics, 2011 Agricultural and Life Sciences Building, Oregon State University, Corvallis, OR 97331-7305, USA.

Our laboratory has reported data suggesting a role for T4 phage gene 32 single-stranded DNA-binding protein in organizing a complex of deoxyribonucleotide-synthesizing enzymes at the replication fork. In this article we examined the effects of gene 32 ablation on the association of these enzymes with DNA-protein complexes. These experiments showed several deoxyribonucleotide-synthesizing enzymes to be present in DNA-protein complexes, with some of these associations being dependent on gene 32 protein. To further understand the role of gp32, we created amber mutations at codons 24 and 204 of gene 32, which encodes a 301-residue protein. We used the newly created mutants along with several experimental approaches—DNA-cellulose chromatography, immunoprecipitation, optical biosensor analysis and glutathione-S-transferase pull-downs—to identify relevant protein-protein and protein-DNA interactions. These experiments identified several proteins whose interactions with DNA depend on the presence of intact gp32, notably thymidylate synthase, dihydrofolate (DHF) reductase, ribonucleotide reductase (RNR) and Escherichia coli nucleoside diphosphate (NDP) kinase, and they also demonstrated direct associations between gp32 and RNR and NDP kinase, but not dCMP hydroxymethylase, deoxyribonucleoside monophosphate kinase, or DHF reductase. Taken together, the results support the hypothesis that the gene 32 protein helps to recruit enzymes of deoxyribonucleoside triphosphates synthesis to DNA replication sites.

PMID: 15720556 [PubMed - in process]

Display

Abstract



Show 20



Sort by



Send to




[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 29 2005 17:30:14



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for

Go Clear

Limits

Preview/Index

History

Clipboard

Details

Display

Abstract

Show

20

Sort by

Send to

All: 1

Review: 0

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: J Biol Chem. 2000 Oct 6;275(40):31496-504.

Related Articles, Links

FREE full-text article at
www.jbc.org

Mutations in the N-terminal cooperativity domain of gene 32 protein alter properties of the T4 DNA replication and recombination systems.

Villemain JL, Ma Y, Giedroc DP, Morrical SW.

Department of Biochemistry and Biophysics, Texas A & M University, College Station, Texas 77843-2128, USA.

The gene 32 protein (gp32) of bacteriophage T4 is the essential single-stranded DNA (ssDNA)-binding protein required for phage DNA replication and recombination. gp32 binds ssDNA with high affinity and cooperativity, forming contiguous clusters that optimally configure the ssDNA for recognition by DNA polymerase or recombination enzymes. The precise roles of gp32 affinity and cooperativity in promoting replication and recombination have yet to be defined, however. Previous work established that the N-terminal "B-domain" of gp32 is essential for cooperativity and that point mutations at Arg (4) and Lys(3) positions have varying and dramatic effects on gp32-ssDNA interactions. Therefore, we examined the effects of six different gp32 B-domain mutants on T4 in vitro systems for DNA synthesis and homologous pairing. We find that the B-domain is essential for gp32's stimulation of these reactions. The stimulatory efficacy of gp32 B-domain mutants generally correlates with the hierarchy of relative ssDNA binding affinities, i.e. wild-type gp32 approximately R4K > K3A approximately R4Q > R4T > R4G gp32-B. However, the functional defect of a particular mutant is often greater than can be explained simply by its ability to saturate the ssDNA at equilibrium, suggesting additional defects in the proper assembly and activity of DNA polymerase and recombinase complexes on ssDNA, which may derive from a decreased lifetime of gp32-ssDNA clusters.

PMID: 10906124 [PubMed - indexed for MEDLINE]

Display Abstract

Show

20

Sort by

Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Mar 29 2005 17:30:14